

Table O-1g. Average Monthly Habitat Conditions for Rush, Lee Vining, Parker, and Walker Creeks under Wet, Normal, and Dry Hydrologic Year-Type Conditions for the 6,410-Ft Alternative (Based on Tennant Method)

Creek	Hydrologic Condition	April	May	June	July	August	September	October	November	December	January	February	March	Mean
Rush	Wet	4	3	2	2	2	4	4	4	4.5	4.5	4.5	4.5	3.6
	Normal	5	3	2	4	5	5	5	5	4	4	4	3	4.1
	Dry	5	4	4	5	4	3	4.5	4.5	4	4	4	3	4.1
Lee Vining	Wet	4	2	2	2	3	5	5	5	4	4	4	3	3.6
	Normal	5	3	2	3	5	4	4.5	4	3	3	3	3	3.5
	Dry	4	4	3	5	3	2	4	3	3	3	3	3	3.3
Parker	Wet	2	4	2	4	5	3	4	3	3	3	3	3	3.2
	Normal	2	4	2	4	5	3	4	3	3	3	3	3	3.2
	Dry	2	4	3	4	5	3	4	3	3	3	3	3	3.3
Walker	Wet	1	2	2	5	3	1	4	4	4	3	3	3	2.9
	Normal	1	2	2	5	3	1	4	4	4	3	3	3	2.9
	Dry	1	2	3	5	3	1	4	4	4	3	3	3	3.0

Habitat conditions are as follows:

- 5 = optimum.
- 4.5 = outstanding.
- 4 = excellent.
- 3 = good.
- 2 = fair.
- 1 = poor.
- 0 = severe degradation.