

Table O-1h. Average Monthly Habitat Conditions for Rush, Lee Vining, Parker, and Walker Creeks under Wet, Normal, and Dry Hydrologic Year-Type Conditions for the No-Diversion Alternative (Based on Tennant Method)

Creek	Hydrologic Condition	April	May	June	July	August	September	October	November	December	January	February	March	Mean
Rush	Wet	4	3	2	2	4	5	5	5	5	5	5	5	4.2
	Normal	5	4	4	4	5	5	5	5	5	5	5	5	4.8
	Dry	5	4	3	3	3	3	4.5	4.5	4.5	4.5	4.5	4.5	4.0
Lee Vining	Wet	4	2	2	2	3	5	5	5	5	4.5	4.5	4.5	3.9
	Normal	5	3	2	3	5	4	4.5	4.5	4.5	4.5	4.5	4.5	4.1
	Dry	4	4	3	5	3	2	4.5	4.5	4	4	4	4	3.8
Parker	Wet	5	3	2	2	2	4	5	4.5	4.5	4.5	4.5	4.5	3.8
	Normal	4	4	2	2	4	5	4.5	4	4	4	4	4	3.8
	Dry	3	5	3	3	4	4	4	3	3	3	3	3	3.4
Walker	Wet	4	2	2	2	3	5	4	4	5	4.5	4.5	4.5	3.7
	Normal	3	4	2	3	5	4	5	5	4.5	4.5	4.5	4.5	4.1
	Dry	1	4	3	5	4	2	4.5	4.5	4	4	4	4	3.7

Habitat conditions are as follows:

- 5 = optimum.
- 4.5 = outstanding.
- 4 = excellent.
- 3 = good.
- 2 = fair.
- 1 = poor.
- 0 = severe degradation.